

## Literatur zur elektrischen Impedanzanalyse

Chertow GM, Lowrie EG, Wilmore DW et al. **Nutritional assessment with bioelectrical impedance analysis in maintenance hemodialysis patients.** J Am Soc Nephrol 2005;6:75-81.

Chumlea W C., Guo S S, Kuczmarski Rj, Flegal K M, Johnson C L, Heymsfield S B, Lukawsk H C, Hubbard Friedl V S. **Body Composition estimates from NHANES III Bioelectrical Impedance data.** Intern. Journ. Of Obesity 2002;26,1596-1609.

De Palo T, Messina G, Edefonti A et al. **Normal values of the bioelectrical impedance vector in childhood and puberty.** Nutrition 2000;16:417-24.

Eisenkolbl J, Kartasurya M, Widhalm K. **Underestimation of percentage fat mass measured by bioelectrical impedance analysis compared to dual energy X-ray absorptiometry method in obese children.** Eur.J.Clin. Nutr. 2001;55:423-9.

Faisy C, Rabbat A, Kouchakji B, Laaban JP. **Bioelectrical impedance analysis in estimating nutritional status and outcome of patients with chronic obstructive pulmonary disease and acute respiratory failure.** Intensive Care Med. 2000;26:518-25.

Fischer H, Lembcke B. **Die Anwendung der Bioelektrischen Impedanz-Analyse (BIA) zur Beurteilung der Körperzusammensetzung und des Ernährungszustandes.** Innere Medizin 18 (1/91);13-17.

Grasso G, Alafaci C, Passalacqua M, Morabito A, Buemi M, Salpietro F M, Tomasello F. **Assessment of Human Brain Water Content by Cerebral Bioelectrical Impedance Analysis: A New Technique and its Application to Cerebral Pathological Conditions.** Neurosurgery 2002;Vol.50 N.5:1064-1074.

Gray D.S, Bray A.G, Gemayel N Kaplan K. **Effect of obesity on bioelectrical impedance.** Am. J. Clin. Nutr. 1989;50:255-260.

Gray D.S. **Changes in bioelectrical impedance during fasting.** Am. J. Clin. Nutr. 1988;48:1184-1187.

Gupta D, Lis CG, Dahlk SL, Vashi PG, Grutsch JF, Lammersfeld CA. **Bioelectrical impedance phase angle as a prognostic indicator in advanced pancreatic cancer.** Br.J.Nutr. 2004;92:957-62.

Gupta D, Lammersfeld CA, Burrows JL et al. **Bioelectrical impedance phase angle in clinical practice: implications for prognosis in advanced colorectal cancer.** Am.J.Clin.Nutr. 2004;80:1634-8.

Heitman B.L. (Dept. of Human Nutrition Copenhagen-Denmark). **Prediction Of Body Water And Fat In Adult Danes From Measurement Of Electrical Impedance: A Validation Study.** International Journal of Obesity (1990);14:789-802.

Kandarina B.J.I. **Die klinische Anwendung der bioelektrischen Impedanzanalyse im Vergleich mit der Anthropometrie zur Bestimmung der Körperzusammensetzung bei Gesunden und Kranken.** Dr. Köster Verlag, Berlin 1995.

Kotler D.P, Buraster S, Wang J, Pierson jr. R.N. **Prediction Of Body Cell Mass, Fat-Free Mass And Total Body Water With Bioelectrical Impedance Analysis: Effects Of Race, Sex And Disease.** Am. J. Clin. Nutr. 1996;64 (suppl.):489S-97S.

K.R, Segal-M, Van Loan-P, Fitzgerald-J.A, Hodgdon-T.B, Van Itallie (Dept. of Pediatric Cardiology-School of Medicine N. Y. W. Human Research Center S. Francisco). **Lean Body Mass Estimation By Bioelectrical, Impedance Analysis: A Four Site Cross-Validation Study.** Am. J. Clin. Nutr. 1988;47:7-14.

Kushner Robert F, Schoeller Dale A, Fjeld Carla R, Danford Lynn (Clinical Nutrition Research Unit, Univ. of Chicago and Washington Univ. School of Medicine). **Is The Impedance Index (HT2/R) Significant In Predicting Total Body Water?** Am. J. Clin. Nutr. 1992;56:835-9.

Kusssmaul B, Döring A, Filipiak B. **Bioelektrische Impedanz-Analyse (BIA) in einer epidemiologischen Studie.** Ernährungs-Umschau 43 (1996) Heft 2;46-48.

Kyle UG, Bosaeus I, De Lorenzo AD et al. **Bioelectrical impedance analysis-part I: review of principles and methods.** Clin.Nutr. 2004;23:1226-43.

Kyle UG, Bosaeus I, De Lorenzo AD et al. **Bioelectrical impedance analysis-part II: utilization in clinical practice.** Clin.Nutr. 2004;23:1430-53.

Lembcke M, Fischer H, Jager R, Polat H, Geier H, Rech M, Staszewski S, Helm E, Caspary W (Div. Of Gastroenterology-Dep. of Medicine, Univ. Wolfgang Goethe-Frankfurt Germany). **Early Changes Of Body Composition In Human Immunodeficiency In Virus-Infected Patients: Tetrapolar Body Impedance Analysis Indicates Significant Malnutrition.** Am. J. Clin. Nutr. 1993; 57:15-19.

Maggiore Q, Nigrelli S, Ciccarelli C, Grimaldi C, Rossi G.A, Michelassi C. **Nutritional And Prognostic Correlates Of Bioimpedance Indexes In Hemodialysis Patients.** Kidney International, Vol. 50 (1996);2103-2108.

Mayfield SR, Uauy R, Waidelich D. **Body composition of low-birth-weight infants determined by using bioelectrical resistance and reactance.** Am J Clin Nutr 2005;54:296-303.

Ott M, Fischer H, Polat H, Helm E.B, Frenz M, Caspary W.F, Lembcke B. **Bioelectrical Impedance Analysis As A Predictor Of Survival In Patients With Human Immunodeficiency Virus Infection.** Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology 9:20-25 1995-Raven Press.

Piccoli A. **Bioelectric impedance vector distribution in peritoneal dialysis patients with different hydration status.** Kidney Int. 2004;65:1050-63.

Piccoli A, Fanos V, Peruzzi L et al. **Reference values of the bioelectrical impedance vector in neonates in the first week after birth.** Nutrition 2002;18:383-7.

Piccoli A, Pillon L, Dumler F. **Impedance Vector Distribution by Sex, Race, Body Mass Index, and Age in the United States: Standard Reference Intervals as Bivariate Z Scores.** Nutrition 2002;18:153-167.

Piccoli A, Pittoni G, Facco E, Favaro E, Pillon L. **Relationship between central venous pressure and bioimpedance vector analysis in critically ill patients.** Crit Care Med. 2000;28:132-7.

Piccoli A, Rossi B, Pillon L, Bucciante G (Institute of Internal Medicine, Division of Nephrology, and Clinical Nutrition Unit, University of Padova). **A New Method For Monitoring Body Fluid Variation By Bioimpedance Analysis: The RXc Graph.** Kidney International, Vol. 46 (1994);534-539.

Pirlich M, Plauth M, Lochs H. **Bioelektrische Impedanzanalyse: Fehlerquellen und methodische Grenzen bei der klinischen Anwendung zur Analyse der Körperzusammensetzung.** Akt. Ernähr.-Med. 24 (1999);81-90.

Pirlich M, Schutz T, Spachos T et al. **Bioelectrical impedance analysis is a useful bedside technique to assess malnutrition in cirrhotic patients with and without ascites.** Hepatology 2000;32:1208-15.

Scharfetter H, Schlager T, Stollberger R, Felsberger R, Hutten H, Hinghofer-Szalkay H. **Assessing abdominal fatness with local bioimpedance analysis: basics and experimental findings.** Int.J.Obes.Relat Metab Disord. 2001;25:502-11.

Schols-A, Dingemans-P, Soeters-E, Wouters (Dept. of Pulmonary Disease and Surgery Univ. of Limburg and Asthma Centre-The Netherlands). **Within-Day Variations Of Bioelectrical Resistance Measurements In Patients With Chronic Obstructive Pulmonary Disease.** Clinical Nutrition (1990) 9:266-271.

Segal, K.R. et. al. **Lean body mass estimation by bioelectrical impedance analysis: a four-site cross-validation study.** Am. J. Clin. Nutr. 1988;47:7-14.

Selberg O, Selberg D. **Norms and correlates of bioimpedance phase angle in healthy human subjects, hospitalized patients, and patients with liver cirrhosis.** Eur.J.Appl.Physiol 2002;86:509-16.

Sergi G, Bussolotto M, Perini P, Calliari I, Cecon V, Scanferla F, Bressan M, Moschini G, Enzi G. **Accuracy Of Bioelectrical Impedance Analysis In Estimation Of Extracellular Space In Healthy Subjects And In Fluid Retention States.** Ann. Nutr. Metab. 1994;38:158-165.

Shime N, Ashida H, Chihara E et al. **Bioelectrical impedance analysis for assessment of severity of illness in pediatric patients after heart surgery.** Crit Care Med. 2002;30:518-20.

Shumei S, Sun, Chumlea W C., Heymsfield Steven B., Lukaski Henry C. et al. **Development of bioelectrical impedance analysis prediction equations for body composition with the use of a multicomponent model for use in epidemiologic surveys.** Am. J. Clin Nutr. 2003;77:331-40.

Shizgal H.M. (Dept. of Surgery-Royal Hospital and Mc Gill Univ. Montreal Canada). **Validation Of The Measurement Of Body Composition From Whole Body Bioelectrical Impedance.** Infusionstherapie 1990;17 (suppl. 3):67-74.

Stroh S. **Methoden zur Erfassung der Körperzusammensetzung.** Ernährungs-Umschau 42 (1995) Heft 3;88-94.

Talluri A, Maggia G. **Bioimpedance Analysis (BIA) In Hemodialysis: Technical Aspects.** International Journal of Artificial Organs / Vol. 18 / No.11, 1995;687-692.

Talluri T, Evangelisti A, Liedtke R.J. **Intra/Extra Cellular Fat Free Mass Spaces Defined By Bioelectrical Reactance And Phase Angle.** IX International Conference on Electrical Bio-Impedance; Heidelberg - Germany, September 1995

Tomczak Jörg. **Körperanalysen: Die bioelektrische Impedanzanalyse BIA.** In: F.I.T. Wissenschaftsmagazin der Deutschen Sporthochschule Köln. 1, Nr. (8.Jahrgang), ALPHA Informationsgesellschaft mbH, 2003;34-40.

Toso S, Piccoli A, Gusella M et al. **Altered tissue electric properties in lung cancer patients as detected by bioelectric impedance vector analysis.** Nutrition 2000;16:120-4.

Toso S, Piccoli A, Gusella M et al. **Bioimpedance vector pattern in cancer patients without disease versus locally advanced or disseminated disease.** Nutrition 2003;19:510-4.

### [Physiologische physikalische Grundlagen der elektrischen Impedanzanalyse](#)

### [Methodenvergleich zwischen Densitometrie, Infrarot-Reflektionsmessung und elektrischer Impedanzanalyse zur Bestimmung der Körperzusammensetzung](#)

### [Impedanzanalyse - Methode - Anwendungsgebiete](#)

### [EUSANA Impedanzanalyse-Messgerät \(BIA\)](#)